

Seminar 4: analytic sets, Weierstrass polynomials.

February 23, 2022

Problem 1. Let $U \subset \mathbb{C}^n$ be an open domain, and let $f : U \rightarrow \mathbb{C}$ be a holomorphic function. Let its zero locus $Z := \{f = 0\}$ be non-empty. Let $Z_{crit} \subset Z$ denote the subset consisting of critical points of the function f with critical value 0. Let the complement $Z \setminus Z_{crit}$ be dense in Z . Show that each point $p \in Z_{crit}$ is a singular point of the zero locus Z : this means that there exists no neighborhood $V = V(p) \subset U$ such that $Z \cap V$ is a submanifold in V .

Problem 2. (Homework) Consider an affine chart \mathbb{C}^n in complex projective space $\mathbb{C}\mathbb{P}^n$. Prove that the closure in $\mathbb{C}\mathbb{P}^n$ of the zero set in \mathbb{C}^n of arbitrary polynomial is an analytic subset in $\mathbb{C}\mathbb{P}^n$. Deduce that the union of the infinity hyperplane and the zero set is an analytic subset in $\mathbb{C}\mathbb{P}^n$.

Problem 3. Prove **Weierstrass Division Theorem.** Let $g(z_1, w) = z_1^d + a_1(w)z_1^{d-1} + \dots + a_d(z_1)$ be a Weierstrass polynomial of degree d in $z_1 \in \mathbb{C}$; here $w \in \mathbb{C}^{n-1}$ (more precisely, we deal with its germ at $0 \in \mathbb{C}^n$): $a_1(0) = \dots = a_d(0) = 0$. For every germ of holomorphic function $f(z_1, w)$ there exist a unique germ of holomorphic function $h(z_1, w)$ and a unique polynomial $p(z_1, w) = a_0(w)z_1^\nu + \dots + a_\nu(w)$ in z_1 , $\nu = \deg_{z_1} p < d$, with coefficients $a_j(w)$ holomorphic at $w = 0$ such that

$$(1) \quad f = hg + p.$$

a) Prove a version for one variable: for every holomorphic function $F(u)$ of one variable u on a closed disk \overline{D}_δ , every $d \in \mathbb{N}$ and every polynomial $g(u)$ of degree d with all the roots contained in D_δ there exist a unique holomorphic function h on D_δ and a unique polynomial $p(u)$ of degree less than d such that

$$F(u) = h(u)g(u) + p(u).$$

b)* Let D_δ and a polydisk $\Delta \subset \mathbb{C}_w^{n-1}$ centered at the origin be such that for every $w \in \Delta$ all the zeros of the polynomial $g(z_1, w)$ in z_1 lie in D_δ , and f is holomorphic on $\overline{D}_\delta \times \Delta$. Prove formula (1) for

$$h(z_1, w) = \frac{1}{2\pi i} \oint_{|u|=\delta} \frac{f(u, w)}{g(u, w)} \frac{du}{u - z_1}.$$

Problem 4.

a) Show that the subset $\{w + e^{\frac{1}{z}} = 0\} \subset \mathbb{C}^2 \setminus \{z = 0\}$ is an analytic subset, but its closure in \mathbb{C}^2 is not an analytic subset in \mathbb{C}^2 .

b) Find the minimal analytic subset in \mathbb{C}^2 that contains the latter closure.